

iv) naphthyl,

v) furanyl, oxazolyl, isoxazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, thiophenyl, thiazolyl, isothiazolyl, pyrrolyl, imidazolyl, pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 3-indoxazinyl, 2-

5 benzoxazolyl, 2- or 3-benzothiophenyl, 2- or 3-benzofuranyl, 2- or 3-indolyl, 2-benzthiazolyl, 2-benzimidazolyl, 3-indazolyl, and

vi) pyridinyl, pyridinyl-N-oxide, pyrazinyl, pyrimidinyl, pyridazinyl, 1-, 3- or 4-isoquinolinyl, 2-, 3- or 4-quinolinyl, 2- or 3-quinoxalinyl, 2- or 4-quinazolinyl,

10 8. The compound of claim 1 wherein  $R^2$ , optionally substituted with  $R^q$ , is selected from the group consisting of phenyl, naphthalenyl, pyridinyl, thiophenyl, benzothiophenyl, furanyl, benzofuranyl, indolyl, indolinyl, isoquinolinyl and quinolinyl.

15 9. The compound of claim 1 wherein  $R^2$  is selected from the group consisting of 4-methyl-phenyl, 2-chloro-phenyl, 3-chloro-phenyl, 4-chloro-phenyl, 3,4-dichloro-phenyl, benzo[1,3]dioxol-5-yl, 2,3-dihydro benzo[1,4]dioxin-6-yl, 4-methoxy-phenyl, phenyl, 4-phenoxy-phenyl, naphthalen-2-yl, pyridin-3-yl, 2-chloro-pyridin-3-yl, pyridin-4-ylmethyl, 4-benzyloxy-phenyl, 4-  
20 dimethylamino-phenyl, 4-bromo-3-methyl-phenyl, 3-methoxy-4-methyl-phenyl, 3-cyclopentyloxy-4-methoxy-phenyl, 4-bromo-2-chloro-phenyl, 4-bromo-phenyl, 3-dimethylamino-phenyl, 4-morpholin-1-yl-phenyl, 4-pyrrolidin-1-yl-phenyl, 4-(N-propylamino)-phenyl, 4-(N-isobutylamino)-phenyl, 4-diethylamino-phenyl, 4-(N-allylamino)-phenyl, 4-(N-isopropylamino)-phenyl, 4-(N-methyl-N-propylamino)-phenyl, 4-(N-methyl-N-isopropylamino)-phenyl, 4-(N-methyl-N-ethylamino)-phenyl, 4-amino-phenyl, 4-(N-methyl-N-propylamino)-2-chloro-phenyl, 4-(N-ethyl-N-methylamino)-2-chloro-phenyl, 4-(pyrrolidin-1-yl)-2-chloro-phenyl, 4-azetidiny-phenyl, 4-(pyrrolidin-2-one-1-yl)-phenyl, 4-bromo-3-methyl-phenyl, 4-chloro-3-methyl-phenyl, 1-methyl-5-indolinyl, 5-indolinyl, 5-  
25 isoquinolinyl, 6-quinolinyl, benzo[1,3]diox-5-yl and 7-methoxy-benzofuran-2-yl.

30 9. The compound of claim 1 wherein  $R^q$  is selected from the group consisting of -OH, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, i-propyl, t-butyl, -OCH<sub>3</sub>, -OCH<sub>2</sub>CH<sub>3</sub>, -OCH(CH<sub>3</sub>)<sub>2</sub>, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, -O-cyclopentyl,